arXiv:1105.1585v2 [hep-th] 20 Jun 2011

Spinor with Schrödinger symmetry and non-relativistic supersymmetry

Hiroshi Yoda[1], Shin'ichi Nojiri[1,2]
[1]*Department of Physics, Nagoya University, Nagoya 464-8602, Japan*
[2]*Kobayashi-Maskawa Institute for the Origin of Particles and the Universe, Nagoya University, Nagoya 464-8602, Japan*

We construct the d dimensional "half" Schrödinger equation, which is a kind of the root of the Schrödinger equation, from the $d+1$ dimensional free Dirac equation. The solution of the "half" Schrödinger equation also satisfies the usual free Schrödinger equation. We also find that the explicit transformation laws of the Schrödinger and the half Schrödinger fields under the Schrödinger symmetry transformation are derived by starting from the Klein-Gordon equation and the Dirac equation in $d+1$ dimensions. We derive the 3 and 4 dimensional super-Schrödinger algebra from the superconformal algebra in 4 and 5 dimensions. The algebra is realized by introducing two complex scalar and one (complex) spinor fields and the explicit transformation properties have been found.

PACS numbers: 11.30.Pb ,11.25.Hf ,03.65.-w

I. INTRODUCTION

Motivated with the original AdS/CFT correspondence [1–3], non-relativistic conformal symmetry [4, 5], especially for the cold atom [6] has attracted some attentions as a correspondence between the gravity in the anti-de Sitter (AdS) background and the non-relativistic conformal field theory [7–12]. The non-relativistic conformal symmetry is the Galilean conformal symmetry containing the Galilei symmetry and conformal symmetry.

The symmetry in the free Schrödinger equation [13–16] , called as the Schrödinger symmetry, is a kind of Galilean conformal symmetry. The algebra of the Schrödinger symmetry in $d-1$ spacial dimensions is the subalgebra of $d+1$ dimensional conformal algebra, which is the algebra of the isometry of $d+2$ dimensional AdS space-time. Then the d dimensional ($d-1$ spacial dimensional) free Schrödinger equation can be derived from the $d+1$ dimensional Klein-Gordon equation by choosing the wave function to be an eigenstate of the momentum of a light-cone direction. In this paper, we define the "half" Schrödinger equation, which is a kind of the root of the Schrödinger equation, from the $d+1$ dimensional free Dirac equation[1]. We find the solution of the "half" Schrödinger equation also satisfies the free Schrödinger equation. Since the Klein-Gordon equation and the Dirac equation have conformal symmetry, the explicit transformation laws of the Schrödinger and the half Schrödinger fields under the Schrödinger symmetry transformation can be derived. Since there are scalar field(s) corresponding to the Klein-Gordon field(s) and spinor field(s) to the Dirac field(s), we may consider the supersymmetry [18–21]. We should note that a non-relativistic supersymmetry for the vector and spinor has been considered in [24]. Starting from the superconformal algebra in 4 and 5 dimensions, we derive the 3 and 4 dimensional super-Schrödinger algebra. The algebra is realized by introducing two complex scalar and one (complex) spinor fields.

II. SCHRÖDINGER SYMMETRY FROM THE KLEIN-GORDON AND DIRAC EQUATIONS

We now show that the d dimensional free Schrödinger equation can be derived from $d+1$ dimensional d'Alembert equation (massless Klein-Gordon equation) by choosing the wave function to be an eigenstate of the momentum of a light-cone direction.

We denote the coordinates in $d+1$ dimensional flat space-time by $\{x^\mu\}$ ($\mu = 0, \cdots d$) and choose the metric tensor by $\eta^{\mu\nu} := \mathrm{diag}(-1, 1, \cdots, 1)$. We now define the light-cone coordinates by

$$x^{\pm} := \frac{1}{\sqrt{2}}(x^0 \pm x^d)\,. \qquad (1)$$

---

[1] The Schrödinger symmetry for the spinor field has been investigated in [17].

Then in terms of the light-cone coordinates, the metric tensor has the following form:

$$\eta^{\mu\nu} = \begin{pmatrix} 0 & -1 & & & \\ -1 & 0 & & & \\ & & 1 & & \\ & & & \ddots & \\ & & & & 1 \end{pmatrix} \tag{2}$$

Now $\mu, \nu = +, -, i = 1, \cdots d-1$. Then, for example, we have $x^{\pm} = -x_{\mp}$ and

$$p_{\pm} := \frac{1}{\sqrt{2}}(p_0 \pm p_d)\,, \quad p^{\pm} = -p_{\mp}\,. \tag{3}$$

Now the relativistic energy-momentum relationship for the massless particle $p^{\mu}p_{\mu} = -2p_+p_- + \mathbf{p}^2 = 0$ can be rewritten if we write

$$p_+ = \mathcal{M}\,, \quad p_- \equiv E\,, \tag{4}$$

as

$$E = \frac{1}{2\mathcal{M}}\mathbf{p}^2\,, \tag{5}$$

which is nothing but the expression of the kinetic energy for the non-relativistic particle with mass $\mathcal{M}$ in d dimensional space ($d+1$ space-time dimensions).

Eq. (5) may suggest that the massless Klein-Gordon equation could be rewritten as the free Schrödinger equation. By using the light-cone coordinates, the Klein-Gordon equation has the following form:

$$(-2\partial_+\partial_- + \partial_i\partial_i)\phi = 0\,. \tag{6}$$

We now choose ϕ to be an eigenstate of ∂_+ as $\partial_+\phi = -i\mathcal{M}\phi$ and we regard a light-cone coordinate x^- as a "time"-coordinate $x^- \equiv t$. Then the Klein-Gordon equation (6) can be rewritten as the free Schrödinger equation:

$$i\partial_t\phi = -\frac{1}{2\mathcal{M}}\triangle\phi\,. \tag{7}$$

Since we assume ϕ to be an eigenstate of ∂_+, ϕ has the following form:

$$\phi(x^+, x^-, x^i) = \mathrm{e}^{-i\mathcal{M}x^+}\tilde{\phi}(x^-, x^i)\,. \tag{8}$$

Later we consider the conformal transformation of the Schrödinger field ϕ, where the transformation of x^+ can be interpreted as a phase of $\tilde{\phi}$.

The free massless Dirac field also satisfies the Klein-Gordon equation. Then by rewriting the Dirac equation by choosing the Dirac field ψ to be an eigenstate of ∂_+ as $\partial_+\psi = -i\mathcal{M}\psi$ and regarding light-cone coordinate x^- as a "time"-coordinate $x^- \equiv t$, we obtain the Schrödinger version of the Dirac equation. We now use the following convention for the γ matrices:

$$\begin{aligned} &\{\gamma^\mu, \gamma^\nu\} = 2\eta^{\mu\nu}\,, \quad \gamma^{\mu\dagger} = \gamma^0\gamma^\mu\gamma^0\,, \quad \Sigma^{\mu\nu} := \frac{1}{4}[\gamma^\mu, \gamma^\nu]\,, \\ &\gamma^5 := i\gamma^0\gamma^1\gamma^2\gamma^3\,, \quad \gamma^{\pm} := \frac{1}{\sqrt{2}}(\gamma^0 \pm \gamma^d) = -\frac{1}{\sqrt{2}}(\gamma_0 \mp \gamma_d) =: -\gamma_{\mp}\,. \end{aligned} \tag{9}$$

Then we obtain

$$\{\gamma^+, \gamma^-\} = -2\,, \quad \{\gamma^{\pm}, \gamma^i\} = 0 \quad (i = 1, 2, \cdots, d-1)\ , \quad (\gamma^+)^2 = (\gamma^-)^2 = 0\,. \tag{10}$$

Then by choosing the Dirac field ψ to be an eigenstate of ∂_+ as $\partial_+\psi = -i\mathcal{M}\psi$, the Dirac equation has the following form:

$$\left(i\gamma^a\partial_a + \mathcal{M}\gamma^+\right)\psi = 0 \quad (a = -, i)\ , \tag{11}$$

which tells ψ satisfies the Schrödinger equation:

$$(2i\mathcal{M}\partial_t + \partial_i\partial_i)\psi = 0\,. \tag{12}$$

Then Eq. (11) is the Schrödinger version of the Dirac equation, which we call the "half Schrödinger equation".

The d-dimensional Schrödinger algebra is given as a subalgebra of the $d+1$ dimensional conformal algebra $SO(d+1,2)$, which is given by

$$\begin{aligned}
&[M_{\mu\nu}, M_{\rho\sigma}] = i(\eta_{\mu\sigma}M_{\nu\rho} - \eta_{\mu\rho}M_{\nu\sigma} - \eta_{\nu\sigma}M_{\mu\rho} + \eta_{\nu\rho}M_{\mu\sigma})\,,\\
&[M_{\mu\nu}, P_\rho] = i(\eta_{\nu\rho}P_\mu - \eta_{\mu\rho}P_\nu)\,, \quad [M_{\mu\nu}, K_\rho] = i(\eta_{\nu\rho}K_\mu - \eta_{\mu\rho}K_\nu)\,,\\
&[D, P_\mu] = -iP_\mu\,, \quad [D, K_\mu] = iK_\mu\,, \quad [K_\mu, P_\nu] = -2i(\eta_{\mu\nu}D + M_{\mu\nu})\,.
\end{aligned} \tag{13}$$

Here $M_{\mu\nu}$ generates the Lorentz boost and the rotation and P_μ generates the translation. Furthermore the operator D generates the dilatation

$$x^\mu \to \mathrm{e}^{-\lambda}x^\mu\,, \tag{14}$$

and K_μ generates the special conformal transformation given by

$$x^\mu \to \frac{x^\mu - c^\mu x^2}{1 - 2c^\mu x_\mu + c^2x^2}\,. \tag{15}$$

The $d-1$ dimensional Schrödinger algebra $\mathfrak{sch}(d-1)$ is the subalgebra of the conformal algebra which does not change the eigenvalue $\mathcal{M}$ of P_+, that is, the algebra given by the generators commuting with P_+.

$$\begin{aligned}
&[J_{ij}, J_{kl}] = i(\delta_{il}J_{jk} - \delta_{ik}J_{jl} - \delta_{jl}J_{ik} + \delta_{jk}J_{il})\,,\\
&[J_{ij}, H] = 0\,, \quad [J_{ij}, E] = 0\,, \quad [J_{ij}, F] = 0\,,\\
&[P_i, P_j] = 0\,, \quad [P_i, H] = 0\,, \quad [P_i, E] = iP_i\,, \quad [P_i, F] = 0\,, \quad [J_{ij}, P_k] = i(\delta_{jk}P_i - \delta_{ik}P_j)\,,\\
&[G_i, G_j] = 0\,, \quad [G_i, H] = iP_i\,, \quad [G_i, E] = iG_i\,, \quad [G_i, F] = 0\,, \quad [J_{ij}, G_k] = i(\delta_{jk}G_i - \delta_{ik}G_j)\,,\\
&[P_i, G_j] = i\mathcal{M}\delta_{ij}\,, \quad [E, H] = -2iH\,, \quad [E, F] = 2iF\,, \quad [F, H] = iE\,.
\end{aligned} \tag{16}$$

Here

$$H = P_-\,, \quad P_i = P_i\,, \quad P_+ = \mathcal{M}\,, \quad J_{ij} = M_{ij}\,, \quad G_i = M_{+i}\,, \quad E = D + M_{-+}\,, \quad F = \frac{1}{2}K_+\,. \tag{17}$$

The generators H, P_i, J_{ij}, G_i, E, and F generate the translation of time, the translation of the spacial coordinates, the spacial rotation, the Galilei boost, the dilatation, and the special conformal transformation, which are explicitly given as the following transformation of the coordinates:

$$\begin{aligned}
&H : t \to t + a\,, \quad P_i : x_i \to x_i + b_i\,, \quad J_{ij} : x_i \to R_{ij}x_j \ \left(R_{ij} \in SO\left(d-1\right)\right)\,, \quad G_i : x_i \to x_i + v_i t\,,\\
&E : t \to \lambda^2 t \text{ and } x_i \to \lambda x_i \quad F : t \to \frac{1}{1+ft}t \text{ and } x_i \to \frac{1}{1+ft}x_i\,.
\end{aligned} \tag{18}$$

In (16), since $\mathcal{M}$ can be regarded as a c number, the commutator of P_i and G_j becomes a central extension. In fact, this algebra is one of the central extended of the Galilean conformal algebra which is constructed by nonrelativistic limit ($c \to \infty$) of conformal algebra. [22]

As well known, the Klein-Gordon equation (6) and the Dirac equation are invariant under the conformal transformation and therefore the Schrödinger equation (7) and the half Schrödinger equation (11) is invariant under the Schrödinger transformation. The invariance of the Klein-Gordon and the Dirac equations and the transformation law of the Klein-Gordon and the Dirac fields in general D dimensions are explicitly given in the appendices.

III. SUPER-SCHRÖDINGER SYMMETRY FROM SUPERCONFORMAL SYMMETRY

Since we have found that both of the Klein-Gordon equation and the Dirac equation have the Schrödinger symmetry and found the explicit transformation laws of the Klein-Gordon and the Dirac fields, we now like to find the transformation connecting the Klein-Gordon field(s) and the Dirac field(s), which is the super-Schrödinger symmetry transformation.

The supersymmetric extension of the conformal symmetry is known as the superconformal symmetry. In this section, for simplicity, we construct $(2+1)$ and $(3+1)$ dimensional super-Schrödinger algebra starting with the $D=4$, $\mathcal{N}=1$ and $D=5$, $\mathcal{N}=2$ superconformal algebra.

A. Construction of $(2+1)$ dimensional super-Schrödinger algebra form the reduction of $D=4$, $\mathcal{N}=1$ superconformal algebra

$D=4$, $\mathcal{N}=1$ Superconformal algebra[23], which is given by, in addition to the conformal algebra $SO(4,2)$ (13),

$$\begin{aligned}
&\{Q_\alpha, \bar{Q}_\beta\} = -\frac{1}{2}i\gamma^\mu_{\alpha\beta}P_\mu\,, \quad \{S_\alpha, \bar{S}_\beta\} = \frac{1}{2}i\gamma^\mu_{\alpha\beta}K_\mu\,, \quad \{Q_\alpha, \bar{S}_\beta\} = -\frac{1}{2}i\delta_{\alpha\beta}D + \frac{1}{2}i\Sigma^{\mu\nu}_{\alpha\beta}M_{\mu\nu} + \gamma^5_{\alpha\beta}A\,,\\
&[Q_\alpha, M_{\mu\nu}] = i(\Sigma_{\mu\nu}Q)_\alpha\,, \quad [S_\alpha, M_{\mu\nu}] = i\left(\Sigma_{\mu\nu}S\right)_\alpha\,, \quad [Q_\alpha, D] = \frac{1}{2}iQ_\alpha\,, \quad [S_\alpha, D] = -\frac{1}{2}iS_\alpha\,,\\
&[Q_\alpha, K_\mu] = -i(\gamma_\mu S)_\alpha\,, \quad [S_\alpha, P_\mu] = i(\gamma_\mu Q)_\alpha\,, \quad [Q_\alpha, A] = -\frac{3}{4}(\gamma^5 Q)_\alpha\,, \quad [S_\alpha, A] = \frac{3}{4}\left(\gamma^5 S\right)_\alpha\,.
\end{aligned} \tag{19}$$

In order to obtain the closed algebra, we define projection operators $\mathcal{P}$ and $\bar{\mathcal{P}}$ as follows

$$\mathcal{P} := -\frac{1}{2}\gamma^-\gamma^+\,, \quad \bar{\mathcal{P}} := -\frac{1}{2}\gamma^+\gamma^-\,, \tag{20}$$

which satisfy the following equations:

$$\begin{aligned}
&\mathcal{P} + \bar{\mathcal{P}} = -\frac{1}{2}\{\gamma^-, \gamma^+\} = 1\,, \quad \mathcal{P}^2 = \frac{1}{4}\gamma^-\gamma^+\gamma^-\gamma^+ = -\frac{1}{2}\gamma^-\gamma^+ = \mathcal{P}\,,\\
&\mathcal{P}\gamma^- = \gamma^-\bar{\mathcal{P}} = \gamma^-\,, \quad \gamma^+\mathcal{P} = \bar{\mathcal{P}}\gamma^- = \gamma^+\,, \quad \mathcal{P}\gamma^+ = \gamma^+\bar{\mathcal{P}} = \gamma^-\mathcal{P} = \bar{\mathcal{P}}\gamma^- = 0\,,\\
&\overline{\eta\mathcal{P}} = -\frac{1}{2}\eta^\dagger\mathcal{P}^\dagger = -\frac{1}{2}\bar{\eta}\gamma^+\gamma^-\,, \quad \bar{\mathcal{P}} = -\mathcal{P}^\dagger := -\frac{1}{2}\gamma^+\gamma^-\,,\\
&\bar{\mathcal{P}}\gamma^a\mathcal{P} = 0 \quad (a = -, 1, \ldots, d-1)\,, \quad \bar{\mathcal{P}}\gamma^+\mathcal{P} = \gamma^+\,.
\end{aligned} \tag{21}$$

If we define

$$T_\alpha := \bar{\mathcal{P}}_{\alpha\beta}S_\beta\,, \quad \bar{T}_\alpha := \bar{S}_\beta\mathcal{P}_{\beta\alpha} = \overline{\bar{\mathcal{P}}_{\alpha\beta}S_\beta}\,, \tag{22}$$

we obtain

$$(\gamma^+T)_\alpha = (\bar{T}\gamma^-)_\alpha = 0\,. \tag{23}$$

Then a closed algebra, which we call the super-Schrödinger algebra, is given by

$$\begin{aligned}
&\{Q_\alpha, \bar{Q}_\beta\} = -\frac{1}{2}i\gamma^-_{\alpha\beta}H - \frac{1}{2}i\gamma^i_{\alpha\beta}P_i - \frac{1}{2}i\gamma^+_{\alpha\beta}\mathcal{M}\,, \quad \{T_\alpha, \bar{T}_\beta\} = i\gamma^+_{\alpha\beta}F\,,\\
&\{Q_\alpha, \bar{T}_\beta\} = \frac{1}{4}i(\gamma^-\gamma^+)_{\alpha\beta}E - \frac{1}{4}i(\Sigma_{ij}\gamma^-\gamma^+)_{\alpha\beta}J_{ij} + \frac{1}{2}i(\gamma^+\gamma_i)_{\alpha\beta}G_i - \frac{1}{2}(\gamma^5\gamma^-\gamma^+)_{\alpha\beta}A\,,\\
&[Q_\alpha, J_{ij}] = i(\Sigma_{ij}Q)_\alpha\,, \quad [T_\alpha, J_{ij}] = i(\Sigma_{ij}T)_\alpha\,, \quad [Q_\alpha, G_i] = -\frac{1}{2}i(\gamma^-\gamma_iQ)_\alpha\,, \quad [T_\alpha, G_i] = 0\,,\\
&[Q_\alpha, E] = -\frac{1}{2}i(\gamma^-\gamma^+Q)_\alpha\,, \quad [T_\alpha, E] = -iT_\alpha\,, \quad [T_\alpha, H] = -i(\gamma^+Q)_\alpha\,, \quad [T_\alpha, P_i] = -\frac{1}{2}i(\gamma^+\gamma^-\gamma_iQ)_\alpha\,,\\
&[T_\alpha, F] = 0\,, \quad [Q_\alpha, F] = \frac{1}{2}i(\gamma^-T)_\alpha\,, \quad [Q_\alpha, A] = -\frac{3}{4}(\gamma^5Q)_\alpha\,, \quad [T_\alpha, A] = \frac{3}{4}(\gamma^5T)_\alpha\,.
\end{aligned} \tag{24}$$

We now consider the following Lagrangian

$$\mathcal{L} = i\bar{\psi}\gamma^\mu\partial_\mu\psi - \frac{1}{2}\sum_{i=i,2}\partial_\mu\phi_i^*\partial^\mu\phi_i\,. \tag{25}$$

The Lagrangian is invariant under the following superconformal transformation:

$$\begin{aligned}
&\delta_Q\phi_1 = -\frac{1}{2}i(\bar{\xi}\psi)\,, \quad \delta_Q\phi_2 = -\frac{1}{2}i(\bar{\xi}\gamma^5\psi)\,, \quad \delta_Q\psi = \frac{1}{2}(i\gamma^-\partial_t + i\gamma^i\partial_i + M\gamma^+)(\xi\phi_1 - \gamma^5\xi\phi_2)\,,\\
&\delta_T\phi_1 = -\frac{1}{2}i(\bar{\eta}\bar{\mathcal{P}}(t\gamma^+ - x_i\gamma_i)\psi)\,, \quad \delta_T\phi_2 = \frac{1}{2}i(\bar{\eta}\bar{\mathcal{P}}\gamma^5(t\gamma^+ - x_i\gamma_i)\psi)\,,\\
&\delta_T\psi = \frac{1}{2}i[(2t\partial_t + x_i\partial_i + 2) + \gamma^+\gamma^i(t\partial_i - i\mathcal{M}x_i) - \Sigma_{ij}(x_i\partial_j - x_j\partial_i)](\phi_1 + \gamma^5\phi_2)\mathcal{P}\eta\,,\\
&\delta_A\phi_1 = -\frac{1}{2}\theta\phi_2\,, \quad \delta_A\phi_2 = \frac{1}{2}\theta\phi_1\,, \quad \delta_A\psi = -\frac{1}{4}\theta\gamma^5\psi\,.
\end{aligned} \tag{26}$$

Here the ξ, η and θ are parameters of the transformation. The transformation (26) is closed on shell.

As we are considering the field theory in 4 dimensions, we may define γ^5 and the chiral projection operators $\mathcal{P}_{L,R}$ by

$$\gamma^5 := i\gamma^0\gamma^1\gamma^2\gamma^3 \,, \quad \mathcal{P}_L := \frac{1}{2}\left(1-\gamma^5\right) \,, \quad \mathcal{P}_R := \frac{1}{2}\left(1+\gamma^5\right) \,. \tag{27}$$

We also define the left and right parts for the Dirac field and also the scalar field by

$$\psi_L := \mathcal{P}_L\psi \,, \quad \psi_R := \mathcal{P}_R\psi \,, \quad \phi_L := \phi_1 - \phi_2 \,, \quad \phi_R := \phi_1 + \phi_2 \,. \tag{28}$$

Then the transformation laws (26) can be rewritten as

$$\begin{aligned}
&\delta_{Q_L}\phi_L = -i(\bar{\xi}\psi_L) \,, \quad \delta_{Q_R}\phi_R = -i(\bar{\xi}\psi_R) \,, \\
&\delta_{Q_R}\psi_L = \frac{1}{2}(i\gamma^-\partial_t + i\gamma_i\partial_i + M\gamma^+)\mathcal{P}_R\xi\phi_L \,, \quad \delta_{Q_L}\psi_R = \frac{1}{2}(i\gamma^-\partial_t + i\gamma_i\partial_i + M\gamma^+)\mathcal{P}_L\xi\phi_R \,, \\
&\delta_{T_R}\phi_L = -ix(\bar{\eta}\bar{\mathcal{P}}(t\gamma^+ - x_i\gamma_i)\psi_L) \,, \quad \delta_{T_L}\phi_R = -ix(\bar{\eta}\bar{\mathcal{P}}(t\gamma^+ - x_i\gamma_i)\psi_R) \,, \\
&\delta_{T_L}\psi_L = \frac{1}{2}i[(2t\partial_t + x_i\partial_i + 2) + \gamma^+\gamma_i(t\partial_i - i\mathcal{M}x_i) - \Sigma_{ij}(x_i\partial_j - x_j\partial_i)]\mathcal{P}\mathcal{P}_L\eta\phi_L \,, \\
&\delta_{T_R}\psi_R = \frac{1}{2}i[(2t\partial_t + x_i\partial_i + 2) + \gamma^+\gamma^i(t\partial_i - i\mathcal{M}x_i) - \Sigma_{ij}(x_i\partial_j - x_j\partial_i)]\mathcal{P}\mathcal{P}_R\eta\phi_R \,, \\
&\delta_{\tilde{A}}\phi_L = -\theta\phi_R \,, \quad \delta_{\tilde{A}}\phi_R = \theta\phi_L \,, \quad \delta_{\tilde{A}}\psi_L = \frac{1}{2}\theta\psi_L \,, \quad \delta_{\tilde{A}}\psi_R = -\frac{1}{2}\theta\psi_R \,.
\end{aligned} \tag{29}$$

Here $\tilde{A} := 2A$. Except the transformation generated by A, the transformations are closed on the left and right parts, respectively.

B. Construction of $(3+1)$ dimensional super-Schrödinger algebra from the reduction of $D=5$, $\mathcal{N}=2$ superconformal algebra

$D=5$, $\mathcal{N}=2$ Superconformal algebra [25][26] is given by,

$$\begin{aligned}
&\{Q_{A\alpha}, \bar{Q}^B_\beta\} = -\frac{1}{2}i\delta^B_A\gamma^\mu_{\alpha\beta}P_\mu \,, \quad \{S_{A\alpha}, \bar{S}^B_\beta\} = \frac{1}{2}i\delta^B_A\gamma^\mu_{\alpha\beta}K_\mu \,, \quad \{Q_{A\alpha}, \bar{S}^B_\beta\} = \delta^B_A\left(\frac{1}{2}i\delta_{\alpha\beta}D - \frac{1}{2}i\Sigma^{\mu\nu}_{\alpha\beta}M_{\mu\nu}\right) + \delta_{\alpha\beta}R^B_A \,, \\
&[Q_{A\alpha}, M_{\mu\nu}] = i(\Sigma_{\mu\nu}Q_A)_\alpha \,, \quad [S_{A\alpha}, M_{\mu\nu}] = i(\Sigma_{\mu\nu}S_A)_\alpha \,, \quad [Q_{A\alpha}, D] = \frac{1}{2}iQ_{A\alpha} \,, \quad [S_{A\alpha}, D] = -\frac{1}{2}iS_{A\alpha} \,, \\
&[Q_{A\alpha}, K_\mu] = -i(\gamma_\mu S_A)_\alpha \,, \quad [S_{A\alpha}, P_\mu] = i(\gamma_\mu Q_A)_\alpha \,, \\
&[Q_{A\alpha}, R_{BC}] = \frac{3}{4}(\varepsilon_{AB}Q_{k\alpha} + \varepsilon_{AC}Q_{j\alpha}) \,, \quad [S_{A\alpha}, R_{BC}] = \frac{3}{4}(\varepsilon_{AB}S_{k\alpha} + \varepsilon_{AC}S_{j\alpha}) \,, \\
&[R_{AB}, R_{CD}] = \frac{3}{4}(\varepsilon_{AC}R_{BD} + \varepsilon_{BD}R_{AC} + \varepsilon_{AD}R_{BC} + \varepsilon_{BC}R_{AD})
\end{aligned} \tag{30}$$

and conformal algebra $SO(5,2)$ (13).

Here fermionic generators with extra indices, $A,B,C... = 1,2$, is $SU(2)$ Majorana spinors and $R_{AB} = R_{BA}$ is $SU(2)$ R-symmetry generators. These indices are raised and lowered with totally antisymmetric tensor ε_{AB} and ε^{AB} ($\varepsilon_{12} = \varepsilon^{12} = 1$), as

$$\zeta^A := \varepsilon^{AB}\zeta_B, \qquad \zeta_A := \zeta^B\varepsilon_{BA} \,, \tag{31}$$

and $SU(2)$ R invariant inner product is defined by contraction of raised and lowered indices,

$$\bar{\zeta}^A_1\zeta_{2A} = \varepsilon^{AB}\bar{\zeta}_{1B}\zeta_{2A} = \varepsilon_{AB}\bar{\zeta}^{1A}\zeta^{2B} \,. \tag{32}$$

In the same way as in the $(2+1)$ dimensional case (22), we reduce this algebra using by projection operators (20), such as

$$
\begin{aligned}
&\{Q_{A\alpha}, \bar{Q}^B_\beta\} = -\frac{1}{2} i\delta^B_A [\gamma^-_{\alpha\beta} H + \gamma_{i\alpha\beta} P_i + \gamma^+_{\alpha\beta}\mathcal{M}] , \quad \{S_{A\alpha}, \bar{T}^B_\beta\} = i\delta^B_A \gamma^+_{\alpha\beta} F , \\
&\{Q_{A\alpha}, \bar{T}^B_\beta\} = -\delta^B_A \left[\frac{1}{4} i(\gamma^-\gamma^+)_{\alpha\beta} E - \frac{1}{4} i(\Sigma_{ij}\gamma^-\gamma^+)_{\alpha\beta} J_{ij} + \frac{1}{2} i(\gamma^+\gamma_i)_{\alpha\beta} G_i\right] - \frac{1}{2}(\gamma^-\gamma^+)_{\alpha\beta} R^B_A , \\
&[Q_{A\alpha}, J_{ij}] = i(\Sigma_{ij} Q_A)_\alpha , \quad [T_{A\alpha}, J_{ij}] = i(\Sigma_{ij} T_A)_\alpha , \quad [Q_{A\alpha}, G_i] = -\frac{1}{2} i(\gamma^-\gamma_i Q_A)_\alpha , \\
&[T_{A\alpha}, G_i] = 0 , \quad [Q_{A\alpha}, E] = -\frac{1}{2} i(\gamma^-\gamma^+ Q_A)_\alpha , \quad [T_{A\alpha}, E] = -iT_{A\alpha} , \quad [T_{A\alpha}, H] = -i(\gamma^+ Q_A)_\alpha , \\
&[T_{A\alpha}, P_i] = -\frac{1}{2} i(\gamma^+\gamma^-\gamma_i Q_A)_\alpha , \quad [T_{A\alpha}, F] = 0 , \quad [Q_{A\alpha}, F] = \frac{1}{2} i(\gamma^- T_A)_\alpha , \\
&[Q_{A\alpha}, R_{BC}] = \frac{3}{4}(\varepsilon_{AB} Q_{C\alpha} + \varepsilon_{AC} Q_{B\alpha}) , \quad [T_{A\alpha}, R_{BC}] = \frac{3}{4}(\varepsilon_{AB} T_{C\alpha} + \varepsilon_{AC} T_{B\alpha}) , \\
&[R_{AB}, R_{CD}] = \frac{3}{4}(\varepsilon_{AC} R_{BD} + \varepsilon_{BD} R_{AC} + \varepsilon_{AD} R_{BC} + \varepsilon_{BC} R_{AD}) .
\end{aligned}
\tag{33}
$$

Thus, the on-shell transformation of hypermultiplet including two scalar fields ϕ^A and one spinor field ψ_α is

$$
\begin{aligned}
&\delta_Q \phi^A = -\frac{1}{2} i(\bar{\xi}^A \psi) , \quad \delta_Q \psi = (i\gamma^- \partial_t + i\gamma_i \partial_i + \mathcal{M}\gamma^+)\xi_A \phi^A , \quad \delta_T \phi^A = -\frac{1}{2} i(\bar{\eta}^A \bar{\mathcal{P}}(t\gamma^+ - x_i\gamma_i)\psi) , \\
&\delta_T \psi = -i[(2t\partial_t + x_i\partial_i + 3) + \gamma^+\gamma_i(t\partial_i - i\mathcal{M}x_i) - \Sigma_{ij}(x_i\partial_j - x_j\partial_i)]\mathcal{P}\eta_A \phi^A , \\
&\delta_R \phi^A = -\frac{3}{2}\theta^A{}_B \Phi^B , \quad \delta_R \psi = 0 .
\end{aligned}
\tag{34}
$$

Here the $SU(2)$ Majorana spinor ξ^A, η^A and 2×2 (anti-)Hermitian traceless matrix $\theta^A{}_B$ are parameters of the transformation. In contrast to the 3 dimensional algebra (except the transformation generated by A), because of R-symmetry invariance of spinor field ψ, the transformations of two scalar fields ϕ^A are not closed.

IV. SUMMARY

In this paper, we have found the "half" Schrödinger equation, which is a kind of the root of the Schrödinger equation, from the $d+1$ dimensional free Dirac equation. Starting from the Klein-Gordon equation and the Dirac equation in $d+1$ dimensions, the explicit transformation laws of the Schrödinger and the half Schrödinger fields under the Schrödinger symmetry transformation have been derived. We have also obtained the 3 and 4 dimensional super-Schrödinger algebra from the superconformal algebra in 4 and 5 dimensions. The algebra is realized by introducing two complex scalar and one (complex) spinor fields and the explicit transformation properties have been found. It could be interesting to find any system which has this kind of supersymmetry.

Acknowledgments

This research has been supported by Global COE Program of Nagoya University (G07) provided by the Ministry of Education, Culture, Sports, Science & Technology and by the JSPS Grant-in-Aid for Scientific Research (S) # 22224003 (SN).

Appendix A: Conformal transformations of scalar and spinor fields

1. Scalar

Here we review on the conformal symmetry in D dimensional space-time and show the invariance of the Klein-Gordon equation and the transformation law of the Klein-Gordon field under the conformal transformation.

We consider the coordinate transformation:

$$
x^\mu = f^\mu\left(\tilde{x}^\nu\right) . \tag{A1}
$$

Under the coordinate transformation, the flat metric in D dimensions

$$ds^2 = \eta_{\mu\nu} dx^\mu dx^\nu \,, \tag{A2}$$

is changed as

$$ds^2 = \eta_{\mu\nu} \frac{\partial f^\mu}{\partial \tilde{x}^\rho} \frac{\partial f^\nu}{\partial \tilde{x}^\sigma} d\tilde{x}^\rho d\tilde{x}^\sigma \,. \tag{A3}$$

When the transformed metric is proportional to $\eta_{\mu\nu}$

$$\eta_{\rho\sigma} \frac{\partial f^\rho}{\partial \tilde{x}^\mu} \frac{\partial f^\sigma}{\partial \tilde{x}^\nu} = C\left(\tilde{x}^\tau\right) \eta_{\mu\nu} \,, \tag{A4}$$

the transformation (A4) is called conformal transformation. Here $C\left(\tilde{x}^\tau\right)$ is a function of the coordinate and determined by multiplying Eq. (A4) with $\eta^{\mu\nu}$ as

$$C\left(\tilde{x}^\tau\right) = \frac{1}{D} \eta^{\mu\nu} \eta_{\rho\sigma} \frac{\partial f^\rho}{\partial \tilde{x}^\mu} \frac{\partial f^\sigma}{\partial \tilde{x}^\nu} \,. \tag{A5}$$

Then substituting (A5) into (A4), we obtain

$$\eta_{\rho\sigma} \frac{\partial f^\rho}{\partial \tilde{x}^\mu} \frac{\partial f^\sigma}{\partial \tilde{x}^\nu} = \frac{1}{D} \eta_{\mu\nu} \eta_{\rho\sigma} \eta_{\alpha\beta} \frac{\partial f^\rho}{\partial \tilde{x}^\alpha} \frac{\partial f^\sigma}{\partial \tilde{x}^\beta} \,, \tag{A6}$$

We call $f^\mu(\tilde{x}^\nu)$ satisfying (A6) as the conformal Killing vector. We now define a matrix $M^\mu_{\ \nu}$ by

$$M^\mu_{\ \nu} \equiv \frac{\partial f^\mu\left(\tilde{x}^\rho\right)}{\partial \tilde{x}^\nu} \,, \tag{A7}$$

Eq. (A6) can be rewritten as

$$\eta_{\mu\nu} M^\mu_{\ \rho} M^\nu_{\ \sigma} = \frac{1}{D} \eta_{\rho\sigma} \eta^{\alpha\beta} \eta_{\mu\nu} M^\mu_{\ \alpha} M^\nu_{\ \beta} \,. \tag{A8}$$

By regarding the quantity in (A8) as a matrix with indexes ρ and σ and considering the determinant, we obtain

$$\eta^{\alpha\beta} \eta_{\mu\nu} M^\mu_{\ \alpha} M^\nu_{\ \beta} = D M^{\frac{2}{D}} \,. \tag{A9}$$

Here M is the determinant of $M^\mu_{\ \nu}$: $M = \det M^\mu_{\ \nu}$.

We now consider the coordinate transformation (A1) for the massless scalar field ϕ:

$$S = \frac{1}{2} \int d^D x \, \eta^{\mu\nu} \partial_\mu \phi \partial_\nu \phi \,. \tag{A10}$$

Then we obtain

$$S = \frac{1}{2} \int d^D \tilde{x} \, M \eta^{\mu\nu} \left(M^{-1}\right)_\mu^{\ \rho} \left(M^{-1}\right)_\nu^{\ \sigma} \partial_\rho \left(M^{\frac{2-D}{2}} \tilde{\phi}\right) \partial_\sigma \left(M^{\frac{2-D}{2}} \tilde{\phi}\right) \,. \tag{A11}$$

Here $\tilde{\phi}$ is defined by

$$\tilde{\phi} = M^{\frac{D-2}{2}} \phi \,. \tag{A12}$$

By using (A8) and (A9), we can rewrite the action (A11) as

$$S = \frac{1}{2} \int d^D \tilde{x} \left\{ \eta^{\mu\nu} \partial_\mu \tilde{\phi} \partial_\nu \tilde{\phi} + 2 \left(\frac{2-D}{2D}\right) \eta^{\mu\nu} \left(M^{-1} \partial_\mu M\right) \tilde{\phi} \partial_\nu \tilde{\phi} + \left(\frac{2-D}{2D}\right)^2 \eta^{\mu\nu} \left(M^{-1} \partial_\mu M\right) \left(M^{-1} \partial_\nu M\right) \tilde{\phi}^2 \right\} \,. \tag{A13}$$

By using the partial integration, we can further rewriting (A13) as

$$S = \frac{1}{2} \int d^D \tilde{x} \left\{ \eta^{\mu\nu} \partial_\mu \tilde{\phi} \partial_\nu \tilde{\phi} - \frac{(D-2)}{4(D-1)} \tilde{\phi}^2 \left(-(D-1) \eta^{\mu\nu} \partial_\mu \partial_\nu \zeta - \frac{(D-2)(D-1)}{4} \eta^{\mu\nu} \partial_\mu \zeta \partial_\nu \zeta \right) \right\} \,. \tag{A14}$$

Here ζ is defined by

$$\zeta \equiv \frac{2}{D} \ln M \,. \tag{A15}$$

Then Eqs. (A3), (A6), and (A7) tell that the metric tensor $g_{\mu\nu}$ is given by

$$g_{\mu\nu} = \mathrm{e}^{\zeta} \eta_{\mu\nu} \,. \tag{A16}$$

For the metric (A16), the scalar curvature R is given by

$$R = \left(- \left(D-1\right) \eta^{\mu\nu} \partial_\mu \partial_\nu \zeta - \frac{\left(D-2\right)\left(D-1\right)}{4} \eta^{\mu\nu} \partial_\mu \zeta \partial_\nu \zeta \right) \mathrm{e}^{\zeta} \,. \tag{A17}$$

Since the metric (A16) is obtained from the flat metric (A2), where the scalar curvature vanishes, the scalar curvature (A17) also vanishes. Then the second term in (A14) vanishes and we find

$$S = \frac{1}{2} \int d^D x \, \eta^{\mu\nu} \partial_\mu \phi \partial_\nu \phi = \frac{1}{2} \int d^D \tilde{x} \, \eta^{\mu\nu} \partial_\mu \tilde{\phi} \partial_\nu \tilde{\phi} \,, \tag{A18}$$

that is, the action (A10) of massless free scalar is invariant under the coordinate transformation (A1) and (A12) where f^μ is the conformal Killing vector satisfying (A6).

2. Spinor

Here show the invariance of the Dirac equation and the transformation law of the Dirac field, that is, spinor field, under the conformal transformation.

In order to investigate the conformal transformation in the spinor field, we consider the infinitesimal transformation

$$x^\mu = \tilde{x}^\mu + \delta x^\mu \,. \tag{A19}$$

Then by defining,

$$m^\mu_{\ \ \nu} \equiv \frac{\partial \delta x^\mu}{\partial \tilde{x}^\nu} \,, \tag{A20}$$

we can rewrite the condition (A8) for the conformal transformation as

$$\frac{1}{2} \left(m_{\mu\nu} + m_{\nu\mu} \right) = \frac{1}{D} \eta_{\mu\nu} m \,, \quad m \equiv m^\mu_{\ \ \mu} \,. \tag{A21}$$

The action of the Dirac fermion is given by

$$S = i \int d^D x \bar{\psi} \gamma^\mu \partial_\mu \psi \,. \tag{A22}$$

We now consider the following infinitesimal transformation:

$$\psi = \left(1 - \frac{D-1}{2D} + \frac{1}{2} m^{\mu\nu} \Sigma_{\mu\nu} \right) \tilde{\psi} \,, \tag{A23}$$

which gives

$$\bar{\psi} = \left(1 - \frac{D-1}{2D} - \frac{1}{2} m^{\mu\nu} \Sigma_{\mu\nu} \right) \tilde{\bar{\psi}} \,, \tag{A24}$$

Then we find

$$S = i \int d^D x \left(\tilde{\bar{\psi}} \gamma^\mu \partial_\mu \tilde{\psi} - m^\nu_{\ \ \mu} \tilde{\bar{\psi}} \gamma^\mu \partial_\nu \tilde{\psi} - m \tilde{\bar{\psi}} \gamma^\mu \partial_\mu \tilde{\psi} - \frac{D-1}{2D} \tilde{\bar{\psi}} \gamma^\mu \tilde{\psi} \partial_\mu m + \frac{1}{2} \tilde{\bar{\psi}} \gamma^\rho \Sigma_{\mu\nu} \tilde{\psi} \partial_\rho m^{\mu\nu} \right) \,. \tag{A25}$$

Since

$$d^D x\,(1-m) = d^D \tilde{x}\,, \quad \tilde{\partial}_\mu \equiv \frac{\partial}{\partial \tilde{x}_\mu} = \frac{\partial}{\partial x_\mu} - m^\nu{}_\mu \frac{\partial}{\partial x_\nu}\,, \tag{A26}$$

we find

$$S = i \int d^D \tilde{x} \left(\tilde{\bar{\psi}} \gamma^\mu \tilde{\partial}_\mu \tilde{\psi} - \frac{D-1}{2D} \tilde{\bar{\psi}} \gamma^\mu \tilde{\psi} \tilde{\partial}_\mu m + \frac{1}{2} \tilde{\bar{\psi}} \gamma^\rho \Sigma_{\mu\nu} \tilde{\psi} \tilde{\partial}_\rho m^{\mu\nu} \right)\,. \tag{A27}$$

In case of rotation (including the Lorentz transformation), translation, and dilatation, $m^{\mu\nu}$ is a constant or vanishes and therefore $\partial_\mu m = \partial_\rho m^{\mu\nu}$, which tells that the action (A22) is invariant under the rotation (including the Lorentz transformation), translation, and dilatation. In case of the special conformal transformation with a infinitesimal parameter c^μ,

$$\delta x = c^\mu \left(\tilde{x}\right)^2 - 2\tilde{x}^\mu \tilde{x}^\rho c_\rho\,, \tag{A28}$$

we find

$$\partial_\mu m = -2Dc_\mu\,, \quad \gamma^\rho \Sigma_{\mu\nu} \partial_\rho m^{\mu\nu} = -2\left(D-1\right) c_\mu \gamma^\mu\,, \tag{A29}$$

and therefore the second and the third te7rms in (A25) cancel with each other and the action (A22) is invariant under the special conformal transformation (A26).

[1] J. M. Maldacena, Adv. Theor. Math. Phys. **2**, 231 (1998) [Int. J. Theor. Phys. **38**, 1113 (1999)] [arXiv:hep-th/9711200].
[2] S. S. Gubser, I. R. Klebanov and A. M. Polyakov, Phys. Lett. B **428**, 105 (1998) [arXiv:hep-th/9802109].
[3] E. Witten, Adv. Theor. Math. Phys. **2**, 253 (1998) [arXiv:hep-th/9802150].
[4] D. T. Son, Phys. Rev. D **78**, 046003 (2008) [arXiv:0804.3972 [hep-th]].
[5] K. Balasubramanian and J. McGreevy, Phys. Rev. Lett. **101**, 061601 (2008) [arXiv:0804.4053 [hep-th]].
[6] Y. Nishida and D. T. Son, Phys. Rev. D **76**, 086004 (2007) [arXiv:0706.3746 [hep-th]].
[7] C. P. Herzog, M. Rangamani and S. F. Ross, JHEP **0811**, 080 (2008) [arXiv:0807.1099 [hep-th]].
[8] J. Maldacena, D. Martelli and Y. Tachikawa, JHEP **0810**, 072 (2008) [arXiv:0807.1100 [hep-th]].
[9] A. Adams, K. Balasubramanian, J. McGreevy JHEP **0811**, 059 (2008) [arXiv:0807.1111 [hep-th]].
[10] W. D. Goldberger, JHEP **0903**, 069 (2009) [arXiv:0806.2867 [hep-th]].
[11] J. L. F. Barbon and C. A. Fuertes, JHEP **0809**, 030 (2008) [arXiv:0806.3244 [hep-th]].
[12] A. Bagchi and R. Gopakumar, JHEP **0907**, 037 (2009) [arXiv:0902.1385 [hep-th]].
[13] C. R. Hagen, Phys. Rev. D **5**, 377 (1972).
[14] U. Niederer, Helv. Phys. Acta **45**, 802 (1972).
[15] M. Henkel, J. Statist. Phys. **75**, 1023 (1994) [arXiv:hep-th/9310081].
[16] T. Mehen, I. W. Stewart and M. B. Wise, Phys. Lett. B **474**, 145 (2000) [arXiv:hep-th/9910025].
[17] C. Duval, P. A. Horvathy, L. Palla, Annals Phys. **249**, 265-297 (1996). [hep-th/9510114].
[18] M. Sakaguchi and K. Yoshida, J. Math. Phys. **49**, 102302 (2008) [arXiv:0805.2661 [hep-th]].
[19] M. Sakaguchi and K. Yoshida, JHEP **0808**, 049 (2008) [arXiv:0806.3612 [hep-th]].
[20] W. Y. Wen, arXiv:0807.0633 [hep-th].
[21] M. Sakaguchi, J. Math. Phys. **51**, 042301 (2010) [arXiv:0905.0188 [hep-th]].
[22] J. Lukierski, P. C. Stichel and W. J. Zakrzewski, Phys. Lett. A**357**, 1 (2006) [arXiv:hep-th/0511259].
[23] M. Kaku, P.K. Townsend, P. van Nieuwenhuizen, Phys. Rev. D **17**, 3179 (1978).
[24] P. A. Horvathy, M. S. Plyushchay, M. Valenzuela, J. Math. Phys. **51**, 092108 (2010). [arXiv:1002.4729 [hep-th]].
[25] T. Fujita and K. Ohashi, Prog. Theor. Phys. **106**, No. 1, 221 (2001) [arXiv:hep-th/0104130].
[26] E. Bergshoeff, S. Cucu, T. de Wit, Jos Gheerardyn, R. Halbersma, S. Vandoren, A. Van Proeyen, JHEP **0210**, 045 (2002) [arXiv:hep-th/0205230].